arXiv:1210.6276v1 [cond-mat.mes-hall] 23 Oct 2012

Nonclassical photon pair production in a voltage-biased Josephson junction

Juha Leppäkangas,[1] Göran Johansson,[1] Michael Marthaler,[2] and Mikael Fogelström[1]

[1] *Microtechnology and Nanoscience, MC2, Chalmers University of Technology, SE-412 96 Göteborg, Sweden*
[2] *Institut für Theoretische Festkörperphysik and DFG-Center for Functional Nanostructures (CFN), Karlsruhe Institute of Technology, D-76128 Karlsruhe, Germany*

We investigate electromagnetic radiation emitted by a small voltage-biased Josephson junction connected to a superconducting transmission line. At frequencies below the well known emission peak at the Josephson frequency ($2eV/h$), extra radiation is triggered by quantum fluctuations in the electromagnetic environment. For weak tunneling couplings and typical ohmic transmission lines, the corresponding photon flux spectrum is symmetric around half the Josephson frequency, indicating that the photons are predominately created in pairs. By establishing an input-output formalism for the microwave field in the transmission line, we give further evidence for this nonclassical photon pair production, demonstrating that it violates the classical Cauchy-Schwarz inequality for two-mode flux cross correlations. In connection to recent experiments, we also consider a stepped transmission line, where resonances increase the signal-to-noise ratio.

PACS numbers: 85.25.Cp,74.50.+r,42.50.Lc

A voltage biased Josephson junction (JJ) produces an oscillating supercurrent, known as the ac Josephson effect [1–3]. In a resistive environment, this current creates electromagnetic (EM) radiation, usually in the microwave regime. The radiation has been studied intensively in literature [3–8], not least from a metrological perspective, since its frequency f is given by the applied voltage V, in the simplest form as $f = 2eV/h$, where e is the electron charge and h is Planck's constant. For usual applications, involving JJs embedded in a low-ohmic environment at moderate temperatures, a classical treatment of the interplay is adequate. Here, the microwave power spectrum has a peak at f broadened by thermal fluctuations in the bias line. At very low temperatures the quantum fluctuations of the resistor set a lower limit to the linewidth [9], through a shot noise in the charge transport [10].

In this Letter, we investigate the microwave field created by very small JJs, where the charge transport takes place through an incoherent sequence of tunneling Cooper-pairs [11–15]. Here, higher-order emission processes can also be detected, without being masked by classical fluctuations. This was recently addressed experimentally [15], by simultaneous measurement of dc current and the power spectrum. It was found that the radiation consists either of a single photon or a pair of photons with the total energy $\sum_i hf_i = 2eV$, corresponding to the energy loss of a single tunneling Cooper-pair. The two-photon emission is triggered by quantum fluctuations, and is of special interest, as this radiation is of nonclassical type. Such production of nonclassically correlated photons has been subject of an active study in various systems taking advantage of non-linearities created by JJs [16–23]. It has a wide interest in the fields of quantum communication [24] and metrology [25–27].

We develop a theoretical framework for perhaps the simplest nonclassical microwave radiation-source based on JJ physics, a dc voltage-biased small JJ. In particular, we establish an input-output theory for the microwave field in the transmission line terminated by a JJ. This provides a straightforward method to calculate any EM-field correlations in the output radiation in the case of weak Josephson coupling. Especially, by calculating photon-flux correlations in the output field, we show that



FIG. 1: (a) A superconducting transmission line terminated by a small Josephson junction. We consider an ohmic transmission line ($Z_0 = Z_1$) and one with a step-like characteristic impedance ($Z_1 \gg Z_0$), supporting modes of a $\lambda/4$-resonator. (b) The equivalent lumped element model. (c) The emitted radiation, either single-photon (I) or two-photon emission (II), originates from the electrostatic energy released by a tunneling Cooper pair. (d) The output photon-flux density vs drive frequency, ω_d, (in units of $I_{\rm c}^2 Z_0/8\hbar\omega_{\rm d}^2$) due to classical Josephson radiation (blue), incoherent Cooper-pair tunneling (red), and thermal radiation (green, $\hbar\omega_{\rm d} = 2E_{\rm J}$, $Z_0 = 50\ \Omega$, $T = 50$ mK). The first two differ in emission at frequencies well below $\omega_{\rm d}$. There the incoherent-Cooper-pair tunneling part is symmetric around the half drive Josephson frequency $\omega_{\rm d}/2$, indicating emission due to pair production of photons.

the radiation is indeed of nonclassical type, as it violates a classical two-mode Cauchy-Schwarz inequality [28–30]. For optimal detection of this, we also consider a stepped transmission line [15], which leads to an increase in the signal-to-noise ratio by enhancing emission at specific frequencies. The estimated rate for creation of photon pairs is of the same order as in the recent two-mode squeezing experiments, made in the context of dynamical Casimir effect [18, 31, 32], showing that the detection of this nonclassical photon production, as a side product of the versatile ac Josephson effect, is in the reach of present technology.

The system we consider consists of a superconducting transmission line (TL) terminated by a small Josephson junction characterized by its capacitance $C_{\rm J}$ and critical current $I_{\rm c}$, see Fig. 1a. The line is dc-voltage biased, with a voltage smaller than the superconducting gap, $eV < 2\Delta$. We focus on the EM radiation in the semi-infinite TL, which we describe by its magnetic flux in a discretized circuit model, Fig. 1b. The TL is characterized by its capacitance C_0 and inductance L_0 per unit length. We now treat the case of a homogenous TL and discuss the stepped impedance case at the end of the paper. In the continuum limit $\delta x \to 0$, the Heisenberg equations of motion for the flux field operator $\Phi(x,t)$ has the form of a Klein-Gordon equation for a massless particle [32], with the traveling-wave solution

$$\begin{aligned}\Phi(x,t) = \sqrt{\frac{\hbar Z_0}{4\pi}} \int_0^\infty \frac{d\omega}{\sqrt{\omega}} \times \\ \left[a_{\rm in}(\omega) e^{-i(-k_\omega x+\omega t)} + a_{\rm out}(\omega) e^{-i(k_\omega x+\omega t)} + {\rm H.c.}\right].\end{aligned} \tag{1}$$

Here $Z_0 = \sqrt{L_0/C_0}$ is the characteristic impedance and $k_\omega = \omega\sqrt{C_0 L_0}$ the wave number. The photon operators $a^\dagger_{\rm in/out}(\omega)$ $[a_{\rm in/out}(\omega)]$ create (annihilate) a photon of frequency ω moving rightwards (in) or leftwards (out) and satisfy the commutation relations $[a_{\rm in(out)}(\omega), a^\dagger_{\rm in(out)}(\omega')] = \delta(\omega-\omega')$. The left- and right-moving part of the field are connected by the boundary condition imposed by the JJ at $x = d$,

$$\begin{aligned}C_{\rm J}\ddot{\Phi}(d,t) \;+\; \frac{1}{L_0}\frac{\partial\Phi(x,t)}{\partial x}|_{x=d} = \\ = -I_{\rm c}\sin\left[2\pi\frac{\Phi(d,t)}{\Phi_0} - \omega_{\rm d} t\right].\end{aligned} \tag{2}$$

Here $\Phi(x=d,t)$ is the magnetic flux across the junction, $\Phi_0 = h/2e$ is the magnetic flux quantum, and $\omega_{\rm d} = 2eV/\hbar$ is the Josephson frequency.

For a steady state (in the Heisenberg picture) one can solve for the output operators $a_{\rm out}(\omega)$ as a function of the input operators $a_{\rm in}(\omega)$. We seek a solution in powers of the critical current $I_{\rm c}$ by multiplying the right-hand side of the boundary condition (2) by ξ and correspondingly write the solution for the outgoing wave in Eq. (1) as $a_{\rm out}(\omega) = \sum_{n=0}^\infty \xi^n a_n(\omega)$. The input field is independent of ξ. The zeroth-order solution (at $x = d$) describes the phase shift given by reflection at the junction capacitance

$$a_0(\omega) = [\chi(\omega)/\chi^*(\omega)]a_{\rm in}(\omega), \tag{3}$$

where $\chi(\omega) = \sqrt{Z_0}/(1 - iZ_0C_{\rm J}\omega)$.

The effect of Cooper-pair tunneling appears in the leading-order solution

$$a_1(\omega) = I_{\rm c}\frac{i\chi(\omega)}{\sqrt{\hbar\omega\pi}}\int_{-\infty}^{\infty} dt\, e^{i\omega t}\sin[\phi_0(t) - \omega_{\rm d} t], \tag{4}$$

where

$$\phi_0(t) = \frac{\sqrt{4\pi\hbar}}{\Phi_0}\int_0^\infty \frac{d\omega}{\sqrt{\omega}}\chi(\omega)a_{\rm in}(\omega)e^{-i\omega t} + {\rm H.c.} \tag{5}$$

is the the zeroth order expression for the phase fluctuation operator at the junction $\phi(t) = (2\pi/\Phi_0)\Phi(d,t)$, giving the dependence on the input.

To second order in $I_{\rm c}$, we find

$$a_2(\omega) = I_{\rm c}^2\frac{i\chi(\omega)}{\sqrt{\hbar\omega\pi}}\int_{-\infty}^{\infty} dt e^{i\omega t}\left[\sin[\phi_0(t) - \omega_{\rm d} t], z(t)\right], \tag{6}$$

where the operator

$$z(t) = \frac{i}{4e}\int_{-\infty}^{\infty} dt'\left[1 + \frac{{\rm Sgn}(t-t')}{e^{\omega_{\rm c}|t-t'|} - 1}\right]\cos[\phi_0(t') - \omega_{\rm d} t'] \tag{7}$$

is a solution to the equation $\phi_1(t) = I_{\rm c}[\phi_0(t), z]$, where $\phi_1(t)$ is the first-order result for the phase difference at the junction. From Eqs. (3), (4) and (6) all correlation functions of the output field can be calculated, to second order in $I_{\rm c}$. This is the main theoretical result in this paper. As a consistency check, we have also verified that these perturbative expressions for the output operators satisfy the canonical commutation relations.

We can now straightforwardly calculate, e.g., the photon-flux density of the output radiation, defined as [33]

$$f(\omega) = \int_0^\infty d\omega' \frac{1}{2\pi}\left\langle a^\dagger_{\rm out}(\omega)a_{\rm out}(\omega')\right\rangle, \tag{8}$$

up to second order in $I_{\rm c}$. The result for typical transmission-line parameters is plotted in Fig. 1d. It can be represented as a sum of two contributions: a part describing thermal radiation $f_{\rm th}(\omega)$ and a part describing radiation originating in incoherent Cooper-pair tunneling $f_{\rm t}(\omega)$, which dominates for $\hbar\omega \gg k_{\rm B}T$. We now introduce the tunnel impedance [13] ${\rm Re}[Z_{\rm t}(\omega)] = |\chi(\omega)|^2 = Z_0/\left[1 + (\omega/\omega_{\rm c})^2\right]$, where the cut-off frequency $\omega_{\rm c} = 1/Z_0C_{\rm J}$ is given by the inverse RC-time of the junction and is much higher than any other relevant frequency in the problem. The expression for the tunnel induced photon flux density now reads

$$f_{\mathrm{t}}(\omega) = \frac{I_{\mathrm{c}}^2 \mathrm{Re}[Z_{\mathrm{t}}(\omega)]}{2\omega}\left[P(\hbar\omega_{\mathrm{d}} - \hbar\omega) + P(-\hbar\omega_{\mathrm{d}} - \hbar\omega)\right], \tag{9}$$

where we have

$$P(E) = \frac{1}{2\pi\hbar}\int_{-\infty}^{\infty} dt e^{J(t)+i\frac{E}{\hbar}t}, \tag{10}$$

and $J(t) = \langle[\phi_0(t) - \phi_0(0)]\phi_0(0)\rangle$ is the phase correlation function at the junction. The function $P(E)$ is the probability density for exchanging a total energy E with the EM environment in a single tunneling event. Here the term $P(\hbar\omega_{\mathrm{d}} - \hbar\omega)$ describes radiation coming from forward-direction Cooper-pair tunneling and the term $P(-\hbar\omega_{\mathrm{d}} - \hbar\omega)$ from the Cooper-pair tunneling against the voltage bias, a process completely suppressed at temperatures $k_B T < 2eV$. The contribution $f_{\mathrm{t}}(\omega)$ agrees with Ref. 15, where the same form is obtained by applying the theory of incoherent Cooper-pair tunneling [13] to calculate the associated photon flux. The simultaneous electric current has then the form $I(V) = (\pi\hbar I_{\mathrm{c}}^2/4e)[P(2eV) - P(-2eV)]$. We note that Eq. (9) for the photon flux is actually valid also when the transmission line has resonances, which then show up as peaks in the $P(E)$ and $Z_{\mathrm{t}}(\omega)$ functions, a case treated at the end of this Letter.

We now focus on the *nonclassical origin* of the photon flux for frequencies $k_B T < \hbar\omega < \hbar\omega_d - k_B T$, where the simple TL environment allows for analytic expressions. At zero temperature, the classical result is $P(E) = \delta(E)$ and the radiation is created at $\omega = \omega_{\mathrm{d}}$ with the total power $I_{\mathrm{c}}^2 \mathrm{Re}[Z_{\mathrm{t}}(\omega_{\mathrm{d}})]/2$. In a quantum treatment of the TL, vacuum fluctuations broadens $P(E)$ towards positive energies and there will be photons emitted also at lower frequencies $\omega < \omega_{\mathrm{d}}$. We use the long-time approximation for the phase correlations [12, 34, 35], $J(t) = -2\rho\left[\ln(\omega_{\mathrm{c}}|t|) + \gamma + i\frac{\pi}{2}\mathrm{sign}(t)\right]$, where $\rho = Z_0/R_Q$ is the ratio between the TL impedance and the quantum resistance $R_Q = h/4e^2$ and γ is the Euler constant. Then, for typical transmission line parameters, $\rho \ll 1$, the resulting finite tail at lower frequencies has the simple expression,

$$f_{\mathrm{t}}(\omega < \omega_{\mathrm{d}}) \approx \frac{\rho I_{\mathrm{c}}^2 Z_0}{\hbar\omega(\omega_{\mathrm{d}} - \omega)}. \tag{11}$$

This part of the photon-flux density is symmetric around the half frequency $\omega_{\mathrm{d}}/2$ indicating that output radiation at these frequencies occurs through pair production of photons symmetrically around $\omega_{\mathrm{d}}/2$, Fig. 1c-d. This symmetry is a central result of this work, and also prevails to the case of resonant transmission line.

Photon pair production in the microwave regime can be achieved also through parametric effects in driven JJ-systems, through conversion of drive photons into the photon pairs [16–23]. In the present case, a static voltage bias is used and each photon pair is instead connected to a tunneling Cooper pair. We also note that the shape of the spectrum in Eq. (11) is inverted compared to the one obtained in the case of the dynamical Casimir effect (DCE)[31] which is $\propto \omega(\omega_d - \omega)$. The difference arises since the Cooper-pair tunneling couples to phase fluctuations which are proportional to $1/\omega$, while for DCE the effective boundary inductance is modulated, which couples to current fluctuations proportional to ω.

To verify that the photons are indeed created in pairs, we need to evaluate the second-order coherence $g^{(2)}(0)$[28] of the out-field, which gives the probability to measure two photons simultaneously normalized to the photon flux. To second order in I_{c}^2 (and for $\rho \ll 1$ and $k_B T \ll \hbar\omega_{\mathrm{d}}$ as above) we find

$$g^{(2)}(0) = \frac{\langle a_{\mathrm{out}}^\dagger a_{\mathrm{out}}^\dagger a_{\mathrm{out}} a_{\mathrm{out}}\rangle}{\langle a_{\mathrm{out}}^\dagger a_{\mathrm{out}}\rangle^2} \approx 1 + \left(\frac{2V}{R_Q I_{\mathrm{c}}}\right)^2, \tag{12}$$

which for low powers ($\propto I_{\mathrm{c}}^2$) can be made arbitrary large, indicating that indeed photons are emitted in pairs.

To test whether the out-field also possess *nonclassical correlations* we calculate the corresponding Cauchy-Schwarz (CS) inequality [36]. In a two-mode case the inequality reads [28, 30]

$$|g_{ab}^{(2)}(0)|^2 \leq g_{aa}^{(2)}(0) g_{bb}^{(2)}(0), \tag{13}$$

where the single-mode second-order coherence is defined as above $g_{ii}^{(2)}(0) = \langle a_i^\dagger a_i^\dagger a_i a_i\rangle/\langle a_i^\dagger a_i\rangle^2$, the photon-flux cross-correlator $g_{ab}^{(2)}(0) = \langle a_a^\dagger a_a a_b^\dagger a_b\rangle/\langle a_a^\dagger a_a\rangle\langle a_b^\dagger a_b\rangle$, and the indices a and b refer to two different modes. We consider two frequency separated modes obtained by detecting the out-field in a small frequency range $\delta\omega$ around ω_a and ω_b [29]. On the left-hand side (LHS) we have the squared probability of simultaneously observing a photon at frequency ω_a and ω_b, which for a classical field is bounded by the right hand side (RHS) product of the probabilities of simultaneous pair detection at the individual frequencies.

With the assumption $\omega_a, \omega_b \gg k_{\mathrm{B}} T$ and in the limit $\delta\omega \to 0$ we get to second-order in I_{c} the equivalent CS inequality for the out-field

$$\begin{aligned} P\left[\hbar(\omega_{\mathrm{d}} - \omega_a - \omega_b)\right]^2 &\leq \\ &P\left[\hbar(\omega_{\mathrm{d}} - 2\omega_a)\right] P\left[\hbar(\omega_{\mathrm{d}} - 2\omega_b)\right]. \end{aligned} \tag{14}$$

We notice that the nonclassicality of the field is determined by the $P(E)$-function only. Now, at zero temperature the probability to absorb energy from the EM environment is zero, i.e., $P(E < 0) = 0$. This implies that the RHS of Eq. (14) is zero for either $\omega_a > \omega_d/2$ or $\omega_b > \omega_d/2$. The LHS is finite for $\omega_a + \omega_b \leq \omega_d$, giving a large regime with nonclassical correlations. For $\omega_a = \omega_b$ both sides are equal, corresponding to degenerate parametric down conversion, and the CS inequality is not violated. This is the leading-order (I_{c}^2) result, when all cor-



FIG. 2: (a) Visualization of violation of the classical CS inequality for flux cross-correlations. Negative values are a sign of nonclassicality. The photon pair production is behind the strong violation nearby the diagonal $\omega_a + \omega_b = \omega_d = 50$ μeV/$\hbar$. We plot here $v/|\mathrm{Max}\{v\}|$, where $v = \mathrm{Sgn(d)}\ln(1 + |\mathrm{d}|)$ and $d/(\mu eV)^2 = -P\left[\hbar(\omega_{\rm d} - \omega_a - \omega_b)\right]^2 + P\left[\hbar(\omega_{\rm d} - 2\omega_a)\right]P\left[\hbar(\omega_{\rm d} - 2\omega_b)\right]$. We have $Z_0 = Z_1 = 50$ Ω and $T = 50$ mK. (b) Photon-flux density due to Cooper-pair tunneling in the neighborhood of the optimal drive point $\omega_{\rm d} = \omega_0 + \omega_1 \approx 141$ μeV/$\hbar$, when $Z_1 \to 10Z_0$ and $E_{\rm J} = (\hbar/2e)I_{\rm c} = 10$ μeV. Here, the photon pairs are mainly emitted at ω_0 and $\omega_1 \approx 3\omega_0$. (c) The *measured* violation of the CS inequality $d \times m$ $[m = -f_{\rm t}(\omega_a)f_{\rm t}(\omega_b)]$ in the diagonal $\omega_{\rm d} = \omega_a + \omega_b$ of (a). (d) The measured violation of the CS inequality corresponding to (c), in diagonal $\omega_{\rm d} = \omega_a + \omega_b = \omega_0 + \omega_1$. The resonance occurs when $\omega_a = \omega_0$.

related photons originate from a single-Cooper-pair tunneling event. Also at finite temperatures, the inequality in Eq. (14) may still be violated, as visualized in Fig. 2 a.

We estimate the photon-flux density in the considered ohmic transmission-line setup to be of the order $f(\omega_{\rm d}/2) \sim 10^{-4} - 10^{-3}$ photons per second per bandwidth, which should be readily detectable with a state-of-the art experimental setup [18]. However, in the measurements of second order flux correlation functions the signal-to-noise ratio decreases substantially. Thus, to verify the nonclassicality of the field it is favorable to use parameters maximizing the product of the photon flux at the detection frequencies ω_a and ω_b, i.e., maximizing $f_{\rm t}(\omega_a)f_{\rm t}(\omega_b)$. Here, we also note that the out-field does not possess any steady-state single mode or two-mode squeezing, in which case nonclassicality could have been shown measuring correlation functions to second order in the amplitude instead of flux[17, 19, 20, 22]. The phase fluctuations across the junction actually influence the field in much the same way as pump phase fluctuations in a parametric amplifier, making the amplitude correlations short-lived.

To increase the photon flux, we consider resonantly shaping the $P(E)$-function using a simple step structure in the characteristic impedance[15], see Fig. 1 a. For $Z_1 \gg Z_0$ this provides modes at frequencies $\omega_n = (1+2n)\omega_0$. Evaluating the output field at $x = 0$, through similar theory as discussed before we obtain that the main results Eqs.(9) and (14) prevail, after the change

$$\chi(\omega) = \frac{2\sqrt{Z_1}e^{-ik^1_\omega d}\sqrt{\frac{Z_1}{Z_0}}}{\mathcal{C}^*(\omega)\left(1 + \frac{Z_1}{Z_0}\right)e^{-2ik^1_\omega d} + \mathcal{C}(\omega)\left(\frac{Z_1}{Z_0} - 1\right)}, \quad (15)$$

where $\mathcal{C}(\omega) = 1 + iZ_1C_{\rm J}\omega$ and $k^1_\omega = \omega\sqrt{C_1L_1}$. This changes the zeroth order phase fluctuation operator in Eq. (5), and thus $P(E)$ through Eq. (10) and the tunnel impedance $\mathrm{Re}[Z_{\rm t}(\omega)] = |\chi(\omega)|^2$.

By aligning the detection frequencies with the first two modes $\omega_a = \omega_0$ and $\omega_b = \omega_1$ and driving at $\omega_{\rm d} = \omega_a + \omega_b$ the photon flux can be enhanced dramatically. In Fig. 2b we plot numerical results for the output photon flux density for a setup with $Z_1 = 10Z_0$, nearby the optimal drive for two-photon emission. The output flux is mostly confined into the two frequencies (of the modes), being even larger than the flux at the drive frequency ω_d. In Fig. 2c-d we then plot the violation of the CS inequality at $\omega_{\rm d} = \omega_a + \omega_b$ for the open-space configuration and the resonant setup, multiplied by the product of the photon flux at the two detection frequencies.

In conclusion, we have derived expressions for the output field operators in a one-dimensional transmission line terminated by a voltage biased Josephson junction, to second order in the critical current of the junction. Using this formalism, we have confirmed the expression for the photon flux density derived in Ref. [15] using a different formalism. Furthermore, by calculating second order flux correlation functions we have established that the photons below the Josephson frequency are mainly emitted in pairs and that the field is indeed nonclassical. Finally, we discuss the possibilities to enhance the photon flux by creating resonances in the transmission line and thus facilitate experimental detection.

We acknowledge financial support from the Swedish Research Council and from the EU through the projects ERC and the PROMISCE.

[1] B. D. Josephson, Phys. Lett. **1**, 251 (1962).

[2] P. W. Anderson and J. M. Rowell, Phys. Rev. Lett. **10**, 230 (1963).

[3] K. K. Likharev, *Dynamics of Josephson Junctions and Circuits* (OPA, Amsterdam, 1986).

[4] A. J. Dahm, A. Denenstein, D. N. Langenberg, W. H. Parker, D. Rogovin, and D. J. Scalapino, Phys. Rev. Lett. **22**, 1416 (1969)

[5] M. J. Stephen, Phys. Rev. Lett. **21**, 1629 (1968); Phys. Rev. **182**, 531 (1969).

[6] P. A. Lee and M. O. Scully, Phys. Rev. B **3**, 769 (1971)

[7] D. Rogovin and D. J. Scalapino, Ann. Phys. **86**, 1 (1974)

[8] Y. Levinson, Phys. Rev. B **67**, 184504 (2003).

[9] R. H. Koch, D. J. Van Harlingen, and J. Clarke Phys. Rev. Lett. **45**, 2132 (1980); Phys. Rev. Lett. **47**, 1216 (1981).

[10] H. Grabert and G.-L. Ingold, Europhys. Lett. **58**, 429 (2002).

[11] S. M. Girvin, L. I. Glazman, M. Jonson, D. R. Penn, and M. D. Stiles, Phys. Rev. Lett. **64**, 3183 (1990)

[12] M. H. Devoret, D. Esteve, H. Grabert, G.-L. Ingold, H. Pothier, and C. Urbina, Phys. Rev. Lett. **64**, 1824 (1990).

[13] G.-L. Ingold and Yu. V. Nazarov, in *Single Charge Tunneling: Coulomb Blockade Phenomena in Nanostructures*, edited by H. Grabert and M. H. Devoret (Plenum, New York, 1992), p.21.

[14] T. Holst, D. Esteve, C. Urbina, and M. H. Devoret, Phys. Rev. Lett. **73**, 3455 (1994).

[15] M. Hofheinz, F. Portier, Q. Baudouin, P. Joyez, D. Vion, P. Bertet, P. Roche, and D. Esteve, Phys. Rev. Lett. **106**, 217005 (2011)

[16] B. Yurke, P. G. Kaminsky, R. E. Miller, E. A. Whittaker, A. D. Smith, A. H. Silver, and R. W. Simon, Phys. Rev. Lett. **60**, 764 (1988)

[17] R. Movshovich, B. Yurke, P. G. Kaminsky, A. D. Smith, A. H. Silver, R. W. Simon, and M. V. Schneider, Phys. Rev. Lett. **65**, 1419 (1990)

[18] C. M. Wilson, G. Johansson, A. Pourkabirian, M. Simoen, J. R. Johansson, T. Duty, F. Nori, and P. Delsing, Nature **479**, 376 (2011).

[19] C. Eichler, D. Bozyigit, C. Lang, M. Baur, L. Steffen, J. M. Fink, S. Filipp, and A. Wallraff, Phys. Rev. Lett. **107**, 113601 (2011).

[20] M. A. Castellanos-Beltran, K. D. Irwin, G. C. Hilton, L. R. Vale, and K. W. Lehnert, Nat. Phys. **4**, 929 (2008).

[21] P. Lähteenmäki, G. S. Paraoanu, J. Hassel, and P. J. Hakonen, arXiv:1111.5608 (2011).

[22] E. Flurin, N. Roch, F. Mallet, M. Devoret, and B. Huard, 1204.0732 (2012).

[23] N. Bergeal, F. Schackert, M. Metcalfe, R. Vijay, V. E. Manucharyan, L. Frunzio, D. E. Prober, R. J. Schoelkopf, S. M. Girvin, and M. H. Devoret, Nature **465**, 64 (2010).

[24] N. Gisin, G. Ribordy, W. Tittel, and H. Zbinden, Rev. Mod. Phys. **74**, 145 (2002)

[25] B. M. Escher, R. L. de Matos Filho, and L. Davidovich, Nature Phys. **7**, 406 (2011)

[26] M. Kira, S. W. Koch, R. P. Smith, A. E. Hunter, and S. T. Cundiff, Nature Phys. **7**, 799 (2011).

[27] P. M. Anisimov, G. M. Raterman, A. Chiruvelli, W. N. Plick, S. D. Huver, H. Lee, and J. P. Dowling, Phys. Rev. Lett. **104**, 103602 (2010).

[28] D. F. Walls and G. J. Milburn, *Quantum Optics* (Springer, Berlin, 2008)

[29] M. P. da Silva, D. Bozyigit, A. Wallraff, and A. Blais, Phys. Rev. A **82**, 043804 (2010).

[30] Y.-q. Li, P. J. Edwards, X. Huang, and Y. Wang, J. Opt. B: Quantum Semiclass. Opt. **2**, 292 (2000).

[31] J. R. Johansson, G. Johansson, C. M. Wilson, and F. Nori, Phys. Rev. Lett. **103**, 147003 (2009)

[32] J. R. Johansson, G. Johansson, C. M. Wilson, and F. Nori, Phys. Rev. A **82**, 052509 (2010).

[33] R. Loudon, *The Quantum Theory of Light* (Oxford University Press, New York, 2010).

[34] G.-L. Ingold and H. Grabert, Phys. Rev. Lett. **83**, 3721 (1999).

[35] H. Grabert, P. Schramm, and G.-L. Ingold, Phys. Rep. **168**, 115 (1988).

[36] A. Miranowicz, M. Bartkowiak, X. Wang, Y. Liu, and F. Nori, Phys. Rev. A **82**, 013824 (2010).

[37] Z. Y. Ou and L. Mandel, Phys. Rev. Lett. **61**, 50 (1988).